

# Biocontrol Using *Trichoderma*

## What is *Trichoderma*?

*Trichoderma* species are fungal organisms that can help farmers grow better crops and have higher income while protecting the soil environment. They are living freely in soil in agricultural and natural environments worldwide.

Commercial products have been developed for plant protection that contain special strains of living *Trichoderma* species, mostly *Trichoderma harzianum*. They have been formulated in a way that farmers and growers can easily apply them in the

field. It is a safe and effective **biocontrol agent** (BCA) that can control a variety of fungal (and bacterial) diseases, such as wilt, damping off, stem rot, and downy mildew.

But not only that! *Trichoderma* is a **yield-booster** that makes crops look healthier, grow bigger, and show better root development.

How is it doing this? *Trichoderma* works best in the soil. It is a biological degrader and competitor of fungal plant pathogens, which has evolved

mechanisms for attacking other fungi in the root zone of a plant. Beyond that it also improves nutrient uptake of plants.



*Trichoderma harzianum*



Farmer in Kampong Chang, Cambodia: She can harvest bigger eggplants because she applied *Trichoderma*.

### How *Trichoderma* works

As a biocontrol agent, the three most important antagonistic processes of *Trichoderma* spp. include:

- \* **Antibiosis:** metabolites that inhibit or kill other microorganisms
- \* **Mycoparasitism:** direct interaction with other fungi through attachment and secretion of lytic enzymes
- \* **Competition:** with pathogens and other fungi for food

When *Trichoderma* spp. colonize plant roots they can stimulate plant growth and protect against infections

### Inside this issue:

Why and When should I Use <i>Trichoderma</i> ?	2
How Effective is <i>Trichoderma</i> ?	3
How to Apply <i>Trichoderma</i> ?	4-5
Biocontrol/IPM and Farm Economics	6
Profitability of Crops	7
References: Facts from Science	8

## HANDS-ON GUIDANCE ON IMPLEMENTING BIOCONTROL AND IPM

## BIOCONTROL USING TRICHODERMA

### WHY AND WHEN SHOULD I USE *TRICHODERMA* ?

You should use *Trichoderma* species when you have problems with soilborne fungal pathogens. Rotting (disintegration of parts of a plant), damping off (rapid death and collapse of very young plants), and wilting (loss of turgidity and drooping of leaves or shoots) is often caused by such soil pathogens. *Trichoderma* spp. is most effectively used as a plant strengthener for seed and seedling diseases, less so against disease of mature crops. Especially, seed treat-

ment (coating) is highly recommendable.

There is also evidence that *Trichoderma*-based products are effective for controlling bacterial (vascular) wilts; for instance bacterial wilts of cucurbits (caused by *Erwinia tracheiphila*, e.g., in cucumber and pumpkin) or bacterial wilts of solanaceous crops (caused by *Ralstonia solanacearum*, e.g., in tomato, eggplant, potato). However, bacterial wilts are best managed with products containing biocontrol agents such as

*Pseudomonas fluorescens* or *Bacillus subtilis*, or by combining these with *Trichoderma* spp.

Yet, there is another reason to use *Trichoderma*: Are you seriously interested to increase your yield? *Trichoderma* has been shown to improve nutrient uptake by plants. Crops just grow bigger and healthier (see page 3).

#### Trichoderma effective against various fungal pathogens

<i>Alternaria alternata</i>	Leaf spot and other diseases
<i>Aspergillus niger</i>	Black mould on fruits
<i>Aspergillus flavus</i>	Post-harvest rot on cereal grains, legumes, and tree nuts
<i>Botrytis cinerea</i>	Grey mould disease on fruits
<i>Colletotrichum</i> spp.	Anthraxnose in various fruits post-harvest
<i>Fusarium</i> spp.	Wide range of fungal infections causing wilting and rotting; e.g., on cereals, banana, vegetables etc.
<i>Phytophthora</i> spp.	Blight in potato and tomato, soybean root rot, and many other crop diseases
<i>Pyricularia oryzae</i>	Rice blast pathogen
<i>Pythium</i> spp.	Damping-off, root rot, stem rot
<i>Rhizoctonia solani</i>	Damping-off in seedlings, black scurf of potatoes, bare patch of cereals, root rot of sugar beet, belly rot of cucumber, sheath blight
<i>Sclerotium</i> (syn. with <i>Typhula</i> ) spp.	Blight of cereals and other crops

### PLANT DISEASES IN PICTURES



Downey mildew of cucumber



Bacterial wilt



*Rhizoctonia* sp. on bean



Rice blast



*Fusarium* wilt

## BIOCONTROL USING TRICHODERMA



### HOW EFFECTIVE IS *TRICHODERMA* ?

ASEAN Sustainable Agrifood Systems (SAS) Project conducted field demonstrations in rice in collaboration with the Cambodian Department of Agriculture, and in vegetables in collaboration with the USAID program HARVEST in Cambodia. Some of the results are shown here. Clearly, using *Trichoderma* spp. as a stand-alone agent, or in combination with compost or manure can significantly boost yields in rice and vegetables. Applying


compost alone could already double, in combination with *Trichoderma* spp. even triple yield in rice! (see graph on the right)

In vegetables, the project witnessed a average yield increase of 17.4% in cucumber, and 11.8% in bitter gourd (see below).

Rice and vegetable crops appeared bigger and healthier under treatment with fungal biocontrol and showed no or reduced disease symptoms.

### “*TRICHODERMA* AND COMPOST ARE YIELD BOOSTERS”

(DATA COURTESY OF DR. SOPHEA KEAN, GDA KHM)



Weight of 10 cucumber seedlings taken randomly from non-treated and *Trichoderma*-treated fields: **31 g versus 51 g**



## BIOCONTROL USING TRICHODERMA

### HOW TO APPLY *TRICHODERMA* ?

Products containing *Trichoderma* species are best used as soil amendments, sometimes also as foliar sprays.

Most importantly, these biocontrol agents work best, if they do not have to compete with already established communities of soil microorganisms. Then, the fungus grows rapidly and can out-compete other microorganisms. Therefore, *Trichoderma* spp. is best introduced during the seedling stage and early growth phase of plants, and in materials (like compost) conducive for growth. Organic fertilizer, cow ma-

nure, or rice-straw compost have all been found to be useful carrier-materials for *Trichoderma* spp.

Temperature and humidity are important factors to consider when applying this BCA in the field. For instance, *Trichoderma harzianum* grows well at 30 C and below, but cannot tolerate more than 36 C. This means application in the field should be carried out towards the evening, best after having irrigated or sprinkled the crops.

Many *Trichoderma* products should be stored cool to

extend their shelf life, which can reach one year if kept in the refrigerator.

Examples for application rates are given in the table below.

The images on the right illustrate in some detail how *Trichoderma* spp. is handled for application in the field or greenhouse.

*“Trichoderma is best used as a soil amendment during the early growth phase of crops”*

#### EXAMPLES FOR APPLICATION RATES OF *TRICHODERMA* SPP.

Product	Formulation (concentration of conidia)	Application rate	Compost application rate
<i>T. harzianum</i> (e.g.; strain produced in Cambodia by GDA)	1 x 10 <sup>7</sup> CFU/g (powder)	<ul style="list-style-type: none"> <li>• With compost: 1 kg powder mixed with 1000 kg compost</li> <li>• As foliar spray: 20 g/20 l water</li> </ul>	RICE: 5–20 tons/ha (= 0.5–2 kg/m <sup>2</sup> ) VEGETABLES: 2 kg/m <sup>2</sup>
<i>T. harzianum</i> (e.g., ‘Trisan’, AppliedChem Thailand; strain AP-001)	2 x 10 <sup>8</sup> CFU/g (powder)	<ul style="list-style-type: none"> <li>• Drenching or spraying on plant: 30–50 g/20 l water</li> </ul>	
<i>T. viride</i> (e.g., ‘Bio-Cure’; T Stanes, India)	2 x 10 <sup>7</sup> CFU/g (powder)	<ul style="list-style-type: none"> <li>• 5–8 kg/ha (= 0.5–0.8 g/m<sup>2</sup>)</li> <li>• With compost: 4 kg in 1000 kg compost or organic fertilizer</li> <li>• Seed dressing: 8 g/kg of seeds along with stickers</li> <li>• Seedling treatment: 20 g/l of water. Roots are dipped into solution for 30 min prior to planting.</li> </ul>	

## BIOCONTROL USING TRICHODERMA

## METHODS OF APPLICATION

Application during the seedling stage:

- 1) Mixing (wetable) powder with water
- 2) Spraying seedlings using knapsack sprayer



During/after transplanting:

- 3) Application at transplanting
- 4) 10 days after transplanting using an injection valve through the drip



- 5) Mixing *Trichoderma* with compost

Stir the *Trichoderma* powder formulation in water until it is completely suspended at a rate of 1 kg per ton of compost.

The compost should be kept moist (not wet), so that the fungal biocontrol agent has optimal growing conditions. Cover the compost heap with a plastic sheet or similar and keep it in a shaded place. If possible, place the compost on a support (e.g., a palette) that allows access of air underneath the heap.

Within 4-5 days of inoculating the compost with *Trichoderma*, successful development will be indicated by a whitish layer on the compost, which is the growing fungal mycelium.



## BIOCONTROL USING TRICHODERMA

# BIOCONTROL/IPM AND FARM ECONOMICS

The main indicators for measuring economic sustainability from a producer's point of view (livelihood) include revenue, costs, and income.

Biocontrol agents (BCA) like *Trichoderma*, and other plant protection inputs are production factors that are accounted for under (variable) costs.

Unfortunately, mis- and overuse of synthetic inputs has created a situation where the farm depletes the productive capacity of its natural and human resources, which is not economically viable.

Ecological, social, and eco-

nomically; each of the three is necessary but only all three together, in harmony and balance, are sustainable.

The economic bottom line places a premium on the present (relative to the future). This is why investment in the (mid to long term) environmental sustainability is not adequately valued.

Only based on this integrity can sustainable farm management maintain and improve competitiveness (sales, inputs, personnel, finance) in order to generate financial surplus and safeguard farm operations in the long term.

BCA help to protect ecological integrity and can replace synthetic pesticides that threaten human and environmental health. Besides, they are also pretty effective (see page 3), securing financial surplus.

It is the goal of Integrated Pest Management (IPM) including BCA to prevent losses due to pests and diseases while minimizing input costs.

To that end, we discuss in the following some strategies how this can be achieved.

*“Trichoderma increases productivity - and profits”*

## FAQ: Pest Management and its Economics in Farm Production

Can pest and disease management increase crop yields?

No. Pest and disease management can only reduce losses, which are caused by pests and pathogens. For lowland rice, estimates indicate an absolute yield loss of 1-2 tons per ha.

If I want a higher yield, what can I do?

Focus your attention on improving soil health and nutrients. Increasing soil organic matter is key to stabilizing and improving soil structure. Low tillage, maintaining soil cover (through crops, crop residues, cover crops, rotations) and applying compost or manure can all contribute to better soil health and, ultimately, higher yields.

*Trichoderma* also has a yield-increasing effect. Thus, it can be used as an amendment for improving soil.

Are insect pests the most important pest?

Not always. In rice, the role of insect pests is generally over-estimated. Rodents and weeds are often more damaging.

Vegetables and fruits can suffer from heavy insect pest attack, but plant diseases are also important. Note: Some insect pests are a problem because of overuse of pesticides (e.g. diamondback moth); here, pesticides are not the solution, but the problem! Pesticide overuse is a major driver for increasing input costs.

Fungal diseases often become more important during the wet season, especially when the cropping pattern (e.g. spacing of plants too narrow) supports fungal growth.

Insect pests are best kept in bay by promoting natural enemies.

When is application of BCA most economic?

Application of BCA is most effective and economic at a moderate to low pest and disease level (targeting larval or early stages of pests) in a preventive rather than reactive manner. This reduces application rates.

What is the economic threshold level?

If the yield that I lose if I do nothing is more worth than the cost of plant protection, then I could apply plant protection measures. However, I have to understand the pests and natural enemies involved, plant tolerance and ability to compensate for damage, other investment opportunities, and expected climatic conditions (weather forecasts).

## PROFITABILITY OF CROPS: FIELD EXPERIENCES FROM SAS

Source: UTT Center for International Trade

**Profitability of Growing Rice**

**COMPARISON AMONG THREE COUNTRIES IN 2012**

	Thailand	Vietnam	Myanmar
Productivity (t/ha)	2.8	5.6	2.6
Inputs (US\$/t)	307	128	224
Revenue (US\$/t)	356	228	334
Profit (US\$/t)	49	100	110

Apart from *Trichoderma* (see page 3) SAS has conducted other field trials and demonstrations in different food and cash crops, including:

- Control of flea beetle using *Bacillus thuringiensis* var. *tenebrionis* and entomopathogenic nematodes in cabbage (Thailand, Brunei Darussalam).
- Control of fruit fly (*Bactrocera* species) in fruit trees by mass trapping using attractants (Indonesia).
- Control of stem borers in rice using sex pheromones (Indonesia).
- Control of coffee berry borer by mass trapping with attractants (Indonesia).

The field experience has shown that with regard to input costs and in relation to crop productivity application of BCA are competitive or even cheaper than synthetic inputs.

Because farmers mis- and overuse synthetic inputs, the alleged advantage of synthetics of being cheaper does not result in higher income of farmers. On the contrary, farmers do experience losses, if the balance between input costs and crop produc-

tivity is not right. Degraded soils and pesticide-induced phytotoxicity further attest to this approach as being unsustainable.

The boxes on this page present examples of simplified profit and loss sheets for rice and vegetables (cabbage). For instance, case studies in cabbage have shown that input costs (including labor) varied between 121% (effectively a loss) and 55% of the revenue (see box below).

In rice, input costs varied between 56% and 86% of the revenue in an inter-country comparison (see above). However, productivity of rice is much lower than that of cabbage or other vegetables. Hence, input costs are much more critical in rice and every measure should be taken to reduce costs.

Because *Trichoderma* not only reduces crop losses due to plant disease but also increases yields (productivity), these fungi are very well suited to improve revenue. If *Trichoderma* is properly applied with compost, it can reduce or substitute expensive synthetic fertilizer.

In view of its soil health promoting properties *Trichoderma* is a truly sustainable approach.

### HOW TO REDUCE INPUT COST?

Pesticides	◇ Reduce number of sprays to 2 or less, considering action at low to medium pest & disease levels. Sometimes control is useful even before planting.
	◇ Protect natural enemies by not spraying if not necessary. For instance, in rice do not spray the first 30 days.
	◇ Use biocontrol instead of synthetic pesticides if possible
	◇ Spraying at pest outbreaks is useless
	◇ Do not listen to arguments aiming at creating fear: Even by doing less you will not lose your entire crop!
	◇ Crop rotation (or mixed cropping) will reduce pests & diseases, diversify your farm production, and increase your income.
Fertilizer and Soil	◇ Determine your soil fertility
	◇ Adjust fertilizer applications according to fertility status
	◇ Use compost or manure to improve soil health
	◇ Reduce tillage and maintain soil cover to increase soil organic matter

### INPUT COST AND PROFIT FOR GROWING VEGETABLES

Item	Cabbage (Case 1)	Cabbage (Case 2)
<b>Productivity (t/ha)</b>	<b>20.3</b>	<b>23.0</b>
<b>Inputs (Costs) (US\$/t)</b>	<b>324 / 246#</b>	<b>156 / 160#</b>
* Labor etc.	* 146	* 108
* Fertilizer	* 90	* 36
* Pesticides /or Bio-control#	* 88 / 9.8#	* 12 / 16#
<b>Revenue (US\$/t)</b>	<b>267</b>	<b>283</b>
<b>Profit/Loss (US\$/t)</b>	<b>-48/21#</b>	<b>127 / 123#</b>



## ASEAN Sustainable Agrifood Systems ( SAS )



50 Department of Agriculture, 4<sup>th</sup> Floor,  
Sittiporn Building, Phaholyothin Road,  
Ladyao, Chatuchak, Bangkok 10900  
Thailand

<http://www.asean-agrifood.org>

### References: Facts from Science

1. Harman, G.E. (2000) Myths and dogmas of biocontrol. Changing perceptions derived from research on *Trichoderma harzianum* T-22. *Plant Disease* 84: 377-393  
(A truly informative and scientifically sound review of the use of *Trichoderma* by one of the founders of research and successful entrepreneurs in this area)
2. Howell, C.R. (2003) Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts. *Plant Disease* 87: 4-10.  
(Discusses the mechanisms that *Trichoderma* spp. employs as known at the time. Still a useful account that prepares for a more recent understanding of the mechanisms)
3. Bhattacharjee, R. & Dey, U. (2014) An overview of fungal and bacterial biopesticides to control plant pathogens/diseases. *African Journal of Microbiology Research* 8: 1749-1762.  
(A recent update that includes some taxonomic information on *Trichoderma* spp. and discusses experiences from known applications including a list of the target diseases of *Trichoderma* spp.)
4. Singh, V., Singh, P.N., Yadav, R.L., Awasthi, S.K., Joshi, B.B., Singh, R.K., Lal, R.J., Duttamajumder, S.K. (2010) Increasing the efficacy of *Trichoderma harzianum* for nutrient uptake and control of red rot in sugarcane. *Journal of Horticulture and Forestry* 2: 66-71.  
(*Trichoderma harzianum* not only controls fungal disease but also improves nutrient uptake of the crop.)
5. Sivan, A., Ucko, O., Chet, I. (1987) Biological control of *Fusarium* crown rot of tomato by *Trichoderma harzianum* under field conditions. *Plant Disease* 71: 587-592.  
(Classical field experiments using *Trichoderma harzianum* for control of fungal disease in tomato.)
6. Kredics, L., Antal, Z., Manczinger, L., Szekeres, A., Kevei, F., Nagy, E. (2003) Influence of environmental parameters on *Trichoderma* strains with biocontrol potential. *Food Technology and Biotechnology* 41: 37-42.  
(A useful review on factors that can influence the performance of *Trichoderma harzianum* such as temperature, pH, pesticides, metal ions, antagonistic soil bacteria, etc.)